



Asian Journal of Phytomedicine and Clinical Research

Journal home page: www.ajpcrjournal.com



EVALUATION OF ANTIBACTERIAL ACTIVITY OF *HIBISCUS SABDARIFFA* LEAF EXTRACT USING DISK DIFFUSION METHOD

Vibha Shree Gupta^{*1}, M. Chandravanshi¹, Wasim Raja², Heena Parwin³

^{1*}Department of Biotechnology, Government Nagarjuna Post Graduate College of Science, Raipur, Chhattisgarh, India.

²Central Laboratory Facility, Chhattisgarh Council of Science and Technology, Raipur, Chhattisgarh, India.

³Department of Biotechnology, MITS School of Biotechnology, Bhubaneswar, Orissa, India.

ABSTRACT

Hibiscus sabdariffa L. is widely used in the preparation of beverages. Active component of this plant exhibit antimicrobial activity, yet little research has been conducted on their possible use in food systems as antimicrobials. The use of this medicinal plant as natural antimicrobial agents is gaining popularity. The *Hibiscus sabdariffa* is widely used for the treatment of diseases. The present study is the continuation of a program aimed at investigation of antimicrobial properties of *Hibiscus sabdariffa* leaf extract. The antibacterial activity was evaluated according to the disk diffusion method by using Gram positive; *B. subtilis*, *S. aureus* and *Enterococcus* and Gram negative; *E. coli*, *Acetobacter*, *citrobacter*, bacteria. This study shows that methenolic leaves extracts of *Hibiscus sabdariffa* Linn inhibit the growth of microorganism's dose dependently. The appear results confirm that the antibacterial activity of *Hibiscus sabdariffa* leaves.

KEYWORDS

Antibacterial, Disk Diffusion method, Bacteria and *Hibiscus sabdariffa*.

Author for Correspondence:

Vibha Shree Gupta,
Department of Biotechnology,
Government Nagarjuna PG College of Science,
Raipur, Chhattisgarh, India.

Email: vibhashreeg07@gmail.com

INTRODUCTION

Hibiscus sabdariffa (Roselle, Ambadi bhaji), a member of the Malvaceae family, is a species of *Hibiscus* native to West Africa, middle East and Indian subtropical. In Indian, Ayurveda medicine has used many herbs such as turmeric possibly as early as 1900 BC (Aggarwal *et al*, 2007)¹. The traditional literatures such as the Rig Veda, and Atharva Veda are some of the earliest available documents detailing the medical knowledge that

October – December

140

formed the basis of the Ayurveda system (Sumner and Judith, 2000)². Some important medicinal plants and minerals used in Ayurveda were later described by ancient Indian herbalist. Many research was carried out on *Hibiscus sabdariffa* L. family Malvaceae calyx ethanol extract in order to ascertain its physiological activity-structure relationship are generally located in areas where it is used in food applications and traditional medicine (Ross, 2003)³. The therapeutic effects of *H. sabdariffa* on lipid metabolism was observed (El-Saadany, et al, 1991⁴, Wang, et al, 2000⁵, Chen, et al, 2003⁶, Carvajal-Zarrabal, et al, 2005⁷, Hirunpanich, et al, 2006⁸, Lin, et al, 2007)⁹. Antihypertensive effects was discovered (Faraji, and Tarkhani, 1999¹⁰, Onyenekwe, et al, 1999¹¹, Odigie, et al., 2003¹², Herrera-Arellano, et al, 2004¹³, Ajay, et al, 2007)¹⁴. The Apoptotic effects of this plant in gastric carcinoma cells was also showed (Chang, et al, 2006¹⁵, Lin, et al, 2007)¹⁶. *H. sabdariffa* L. leaf has also been subjected to this type of scientific studies, particularly leaf ethanolic extracts, which have been found to influence lipid metabolism (Essa, et al, 2009)¹⁷. *Hibiscus sabdariffa* calyx extract contains hibiscus acid, or (+) -hydroxycitric acid, known as (+) -HCA (Ibnusaad and Thomas, 2003¹⁸, Jena, et al, 2002)¹⁹ which was useful in several diseases. It has been proposed as an anti-obesity agent (Sullivan, et al, 1974²⁰, Heymsfield, et al, 1998²¹, Hayamizu, et al, 2003)²². The powder made of dried crushed leaves is called ambadi bhaji. The antioxidants including flavonoids, gossypetine, hibiscetine and sabdaretine was present in plant extract. This plant extract was also rich in riboflavin, ascorbic acid, niacin, carotene, calcium, and iron that are nutritionally important. *Hibiscus sabdariffa* L (HS) (Malvaceae), a common local drink popularly known as zobo in Nigeria and medicinal herb, is used in folk medicine in the treatment of hypertension (Wang et al, 2000⁵, Odigie et al, 2003¹², Olaleye and Akindahunsi, 2005)²³. Therefore, we have planned out the antibacterial activity of *Hibiscus sabdariffa* leaf extract on different type of bacteria.

MATERIAL AND METHODS

Plant Materials

The *Hibiscus sabdariffa* (*Ambadi bhaji*) leaves were collected from local herbal garden of Raipur, Chhattisgarh and dried for few days in shade, which were then powdered and preserved in airtight bottles for further studies.

Extract Preparation

Hibiscus sabdariffa L. Leaves (20g) was extracted in 50% of methanol and Millipore water solvent the supernatant was collected and concentrated in water bath at 40-50 C .The dried powder was kept in air tied box.

Microorganisms

The tested microorganisms included the Gram positive bacteria; *Bacillus subtilis*, *Staphylococcus aureus*, (ATCC 25923) *Enterococcus* and Gram negative bacteria; *Acetobacter*, *Citrobacter*, *Escherichia coli*. These bacteria's strains were procured from National Chemical Laboratory (NCL), Pune, India. The bacteria were grown in the nutrient broth at 37° C and maintained on nutrient agar slants at 4° C.

Antibacterial Assay

Antibacterial activity of *Hibiscus sabdariffa* L. Leaf extract was determined by agar disk diffusion method (Nair, et al, 2005)²⁴ at four concentrations i.e., 100, 75, 50 and 25mg/ml. Muller Hinton agar was prepared according to the manufacturer's instructions and the plates were seeded with appropriate microorganisms (*Bacillus subtilis*, *Staphylococcus aureus*, *Enterococcus*, *Acetobacter*, *Citrobacter* and *Escherichia coli*). Discs of 6 mm diameter were prepared from Whatmann filter paper No. 24 and sterilized. The discs were then impregnated with the extracts and solvent DMSO. Antibiotics for Gram positive (NX – Norfloxacin, OF- Ofloxacin, E-Erythromycin, CFM- Cefixime) and Gram Negative (NX–Norfloxacin, OF- Ofloxacin, E-Erythromycin, CFM- Cefixime).Bacteria were used as standard. The plates were incubate at 37° C for 24 hrs and the zones of inhibition were measured with a measuring scale. Above experiment was carried out in triplicate for their confirmation.

RESULTS AND DISCUSSIONS

The results showed that the extracts of *Hibiscus sabdariffa* L. has a concentration dependent antibacterial activity with more sensitivity for gram positive bacteria than gram negative bacteria used in the extracts of *Hibiscus sabdariffa* showed considerable antibacterial activity at all the four concentrations 100, 75, 50, 25 mg/ml (Table No.1). (Table No.2) shows the sensitivity of the tested bacteria to the standard antibiotics.

The medicinal value of plants lies in some chemical substances that have a definite physiological function in the human body. Some different phytochemicals have been found to possess a wide range of medicinal properties, which may help in protection against various diseases. Mainly, alkaloids protect against chronic diseases; saponins protect against hypercholesterolemia and steroids and triterpenoides show the analgesic properties (Geetha *et al*, 2014)²⁵. Some previous findings revealed that these plant present phenolic compounds including flavonoids are potent antioxidants with reported antimutagenic and anticarcinogenic effects. Tannin contributes various medicinal properties such as antimicrobial, anti-inflammatory and astringent activity (Geetha *et al*, 2014)²⁵.

Through the ages man has learnt to take advantage of the many resources placed at his disposal by nature to meet his essential needs in all fields. As important reserves and sources of abundance, natural resources are indispensable for socio-economic development. Many plants of the Malvaceae family have been found to posses antimicrobial properties against several microorganism of clinical importance. In this study, we reported the antibacterial effect of *Hibiscus sabdariffa* extract.

Our result for antibacterial activity of plant *Hibiscus sabdariffa* shows that antibacterial activity of 50% Methanolic extract of *Hibiscus sabdariffa* against different selected bacteria. The above observations suggests that different concentration (25%, 50%, 75%, 100%) were having good antibacterial activity against Acetobacter,

Citrobacter, E.coli ,Bacilli Subtilis, Staphylococcus aures and Enterococcus. Thus the extract is showed varying activity against all organisms. On comparing the zone of inhibition of extract to that of standard antibiotics extract showed better activity than Ciprofloxacin (CIP), Doripenem (DOR), Ofloxacin (OF), Maxifloxacin (OM) in these conditions. The development of antibiotic resistance has become a global public health challenge which is causing in effectiveness of anti-bacterial agents leading to increase in diseases and death rate. Therefore this study intends to investigate the antibacterial action of *Hibiscus sabdariffa* (Ambadi bhaji) extract against multidrug pathogens.

The antibacterial activity obtained in this study, coincides with those reported by other researchers (Mei-Chi and Che-Yi, 2008)²⁶. The antibacterial activity of selected plant may be attributed to the phytochemical compounds it contains especially those of phenolic nature such as gallic acid, quercetin, rutin and luteolin and analogues which were detected therein as previously reported in literature (El-kamali and Moneer, 2006²⁷, Alaa, 2012)²⁸.

The antibacterial activity of *Hibiscus sabdariffa* calyx extract provide valuable information and highlight the potentiality of this plant in drug development as a candidate source of antibacterial agents being safe and edible. Some more investigation is required to ascertain that the in vitro results are attainable in vivo. Moreover, this study strongly supports the ethnomedical uses of *Hibiscus sabdariffa* calyx in treatment of cough, abscess and biliary conditions which are caused mainly by microbial infections (Morton, 1987)²⁹. It can be reported that *Hibiscus sabdariffa* calyx could be a potential source of antibacterial agents so further studies should be conducted to identify the active constituents responsible for this activity.

Table No.1: Antibacterial activity of Hibiscus sabdariffa Leaves extract against tested bacteria

Test sample concentration in (mg/ml)	Name of the Microorganism (Inhibition Zone in mm)					
	Gram Positive			Gram Negative		
	<i>Bacillus subtilis</i>	<i>S.aureus</i>	<i>Enterococcus</i>	<i>Acinetobacter</i>	<i>Citrobacter</i>	<i>E.coli</i>
100%	10	13	14	10	0.6	13
75%	0.8	15	11	0.9	0.0	14
50%	10	16	12	11	0.8	12
25%	11	12	11	0.8	0.8	10

Table No.2: The study of anti-bacterial activities of standard antibiotics using disk diffusion method

S.No	Name of Bacteria	Zone of Inhibition (In MM)			
		Gram Positive (+)	NX	OF	E
1	<i>Bacillus subtilis</i>	37.00	32.00	15.00	28.00
	<i>S. aureus</i>	34.00	31.00	15.00	09.00
	<i>Enterococcus</i>	39.00	36.00	20.00	35.00
	Gram Negative (-)				
2	<i>Acinetobacter</i>	15.00	09.00	09.00	09.00
	<i>Citrobacter</i>	37.00	30.00	15.00	28.00
	<i>E.coli</i>	34.00	30.00	18.00	21.00

SUMMARY AND CONCLUSION

Hibiscus sabdariffa is an ideal crop for developing countries as it is relatively easy to grow, can be grown as part of multi-cropping systems and can be used as food and fiber. In China the seeds are used for their oil and the plant is used for its medicinal properties, while in West Africa the leaves and powdered seeds are used in meals. It is also used in the pharmaceutical and food industries. Our observations suggest that different concentration were having good antibacterial activity against different bacteria's. Thus the extract is showed varying activity against all organisms.

Experimental data suggested that antibacterial study was done which shows a better antibacterial activity against all the six test gram-positive and gram-negative bacteria species used and shown antibacterial susceptibility to extracts with clear zone of inhibition. So, in future it can be used as an alternate to synthetic antibiotics. Much effort has needed to increase plant extract as a dietary supplement in food to resist the human pathogenic bacterial disease. This work has revealed further potentials of this plant in the area antimicrobial agent. As a result of the high antimicrobial activity, the extract of *Hibiscus sabdariffa* would be considered a safe antimicrobial agent

ACKNOWLEDGEMENT

The authors wish to express their sincere gratitude to Department of Biotechnology, Government Nagarjuna Post Graduate College of Science, Raipur, Chhattisgarh, India for providing necessary facilities to carry out this research work.

CONFLICT OF INTEREST

We declare that we have no conflict of interest.

BIBLIOGRAPHY

- Aggarwal B B, Sundaram C, Malani N, Ichikawa H. Curcumin: the Indian solid gold, *Adv. Exp. Med. Biol. Advances in Experimental Medicine and Biology*, 595, 2007, 1-75.
- Sumner Judith. The Natural History of Medicinal Plants, *Timber Press*, 2000, 16-17
- Ross. Medicinal Plants of the World Vol I: *Chemical Constituents, Traditional and Modern Uses*, *Humana Press*, Totowa, NJ, USA, 2nd Edition, 10, 2003, 491.
- El-Saadany M Z, Sitohy S M, Labib and El-Massry R A. Biochemical dynamics and hypocholesterolemic action of *Hibiscus*

- sabdariffa* (Karkade), *Die Nahrung*, 35(6), 1991, 567-576.
- 5. Wang C J, Wang J M, Lin W L, Chu C Y, Chou F P, Tseng T H. Protective effect of Hibiscus anthocyanins against tert – butyl hydroperoxide – induced hepatic toxicity in rats, *Food Chem. Toxicol.*, 38(5), 2000, 411-416.
 - 6. Chen C C, Hsu J D, Wang S F. *Hibiscus sabdariffa* extract inhibits the development of atherosclerosis in cholesterol-fed rabbits, *Journal of Agricultural and Food Chemistry*, 51(18), 2003, 5472-5477.
 - 7. Carvajal-Zarrabal S M, Waliszewski D M, Barradas- Dermitz. The consumption of *Hibiscus sabdariffa* dried calyx ethanolic extract reduced lipid profile in rats, *Plant Foods for Human Nutrition*, 60(4), 2005, 153-159.
 - 8. Hirunpanich V A, Utaipat N P, Morales I. Hypocholesterolemic and antioxidant effects of aqueous extracts from the dried calyx of *Hibiscus sabdariffa* L. in hypercholesterolemic rats, *Journal of Ethnopharmacology*, 103(2), 2006, 252-260.
 - 9. Lin H H, Chen J H, Kuo W H and Wang C J. Chemo preventive properties of *Hibiscus sabdariffa* L. on human gastric carcinoma cells through apoptosis induction and JNK/p38 MAPK signaling activation, *Chemico-Biological Interactions*, 165(1), 2007, 59-75.
 - 10. Faraji M and Haji Tarkhani A H. The effect of sour tea (*Hibiscus sabdariffa*) on essential hypertension, *Journal of Ethnopharmacology*, 65(3), 1999, 231-236.
 - 11. Onyenekwe P C, Ajani E O, Ameh D A and Gamaniel K S. Antihypertensive effect of roselle (*Hibiscus sabdariffa*) calyx infusion in spontaneously hypertensive rats and a comparison of its toxicity with that in wistar rats, *Cell Biochemistry and Function*, 17(3), 1999, 199-206.
 - 12. Odigie I P, Ettarh R R, Adigun S A. Chronic administration of aqueous extract of *Hibiscus sabdariffa* attenuates hypertension and reverses cardiac hypertrophy in 2K-1C hypertensive rats, *J. Ethnopharmacol.*, 86(2-3), 2003, 181-185.
 - 13. Herrera-Arellano S, Flores-Romero M, Chávez-Soto A and Tortoriello J. Effectiveness and tolerability of a standardized extract from *Hibiscus sabdariffa* in patients with mild to moderate hypertension: a controlled and randomized clinical trial, *Phytomedicine*, 11(5), 2004, 375-382.
 - 14. Ajay M, Chai H J, Mustafa A M, Gilani A H and Mustafa M R. Mechanisms of the anti-hypertensive effect of *Hibiscus sabdariffa* L. calyces, *Journal of Ethnopharmacology*, 109(3), 2007, 388-393.
 - 15. Chang Y C, Huang K X, Huang A C, Ho Y C and Wang C J. *Hibiscus* anthocyanins-rich extract inhibited LDL oxidation and ox LDL-mediated macrophages apoptosis, *Food and Chemical Toxicology*, 44(7), 2006, 1015-1023.
 - 16. Lin T L, Lin H H, Chen C C, Lin M C, Chou M C and Wang C J. *Hibiscus sabdariffa* extract reduces serum cholesterol in men and women, *Nutrition Research*, 27(3), 2007, 140-145.
 - 17. Essa M M and Subramanian P. Influence of *Hibiscus sabdariffa* on the rhythmic alterations of liver markers in experimental hyperammonemic rats, *Biological Rhythm Research*, 40(3), 2009, 273-278.
 - 18. Ibnusaud and Thomas G. Biologically interesting chiral 3,4-disubstituted pyrrolidines from optically active hydroxycitric acid lactones, *Tetrahedron Letters*, 44(6), 2003, 1247-1249.
 - 19. Jena S, Jayaprakasha G K, Singh R P and Sakariah K K. Chemistry and biochemistry of (-)-hydroxycitric acid from Garcinia, *Journal of Agricultural and Food Chemistry*, 50(1), 2002, 10-22.
 - 20. Sullivan C J, Triscari and Hamilton J G. Effect of (-) hydroxycitrate upon the

- accumulation of lipid in the rat: I. Lipogenesis, *Lipids*, 9(2), 1974, 121-128.
21. Heymsfield S B, Allison D B, Vasselli J R, Pietrobelli A, Greenfield D and Nuñez C. "Garcinia cambogia (hydroxycitric acid) as a potential antiobesity agent: a randomized controlled trial, *Journal of the American Medical Association*, 280(18), 1998, 1596-1600.
22. Hayamizu K, Ishii Y, Kaneko I. Effect of *Garcinia cambogia* (hydroxycitric acid) on visceral fat accumulation: a double-blind, randomized, placebo-controlled trial, *Current Therapeutic Research*, 64(8), 2003, 551-567.
23. Olaleye M T, Akindahunsi A A. Hypotensive activity of methanolic extract of the calyces of *Hibiscus sabdariffa* L on Normotensive rats. In Recent Progress in Medicinal plants Plant Bioactives in Traditional Medicine, *Stadium press LLC*, U.S.A, 12(9), 2005, 283-288.
24. Nair P N, Henry S, Cano V, Vera J. Microbial status of apical root canal system of human mandibular first molars with primary apical periodontitis after one-visit endodontic treatment, *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*, 99(2), 2005, 231-52.
25. Geetha T S and Geetha N. Phytochemical Screening, Quantitative Analysis of Primary and Secondary Metabolites of *Cymbopogon citratus* (DC) stapf. leaves from Kodaikanal hills, Tamilnadu, *International Journal of Pharm Tech Research*, 6(2), 2014, 521-529.
26. Mei-Chi Y and Che-Yi C. Anticampylobacter, anti-aerobic and antioxidative effects of roselle calyx extract and protocatechuic acid in ground beef, *International Journal of Food Microbiology*, 127(1-2), 2008, 73-77.
27. El-kamali H and Moneer M. Antibacterial activity of *Hibiscus sabdariffa* and *Acacia seyal* against upper respiratory tract pathogens, *Sudan Medical Journal*, 1(2), 2006, 121-126.
28. Alaa G. Antioxidant and antibacterial activities of *Hibiscus sabdariffa* L.Extracts, *African Journal of Food Science*, 6(21), 2012, 506-511.
29. Morton J F. Roselle. In: Fruits of Warm Climate (Dowling CF, ed). Media, Inc., *Greensboro, NC*, 1987, 281-286.
30. Essa M M, Subramanian P, Suthakar G. Influence of *Hibiscus sabdariffa* (Gongura) on the levels of circulatory lipid peroxidation products and liver marker enzymes in experimental hyperammonemia, *Journal of Applied Biomedicine*, 4(1), 2006, 53-58.

Please cite this article in press as: Vibha Shree Gupta *et al.* Evaluation of antibacterial activity of *hibiscus sabdariffa* leaf extract using disk diffusion method, *Asian Journal of Phytomedicine and Clinical Research*, 6(4), 2018, 140-145.